

## Key References for Module 2

- Bagchi, S., & M.E. Ritchie. (2010). Herbivore effects on above-and belowground plant production and soil nitrogen availability in the Trans-Himalayan shrub-steppes. *Oecologia*, 164, 1075-1082.
- Carbone, C., & Gittleman, J. L. (2002). A common rule for the scaling of carnivore density. *Science*, 295(5563), 2273-2276.
- Forsyth, D. M., & Hickling G. J. (1997). An improved technique for indexing abundance of Himalayan thar. *New Zealand journal of ecology*, 97-101.
- Johansson, Ö., McCarthy, T., Samelius, G., Andrén, H., Tumursukh, L., & Mishra, C. (2015). Snow leopard predation in a livestock dominated landscape in Mongolia. *Biological Conservation*, 184, 251-258.
- Rowcliffe, J. M., Field, J., Turvey, S. T., & Carbone, C. (2008). Estimating animal density using camera traps without the need for individual recognition. *Journal of Applied Ecology*, 1228-1236.
- Singh, N.J., & Milner-Gulland, E.J. (2010). Monitoring ungulates in Central Asia: current constraints and future potential. *Oryx*, doi: :10.1017/S0030605310000839
- Suryawanshi, K. R., Bhatnagar, Y. V., & Mishra, C. (2012). Standardizing the double-observer survey method for estimating mountain ungulate prey of the endangered snow leopard. *Oecologia*, 169(3), 581-590.
- Suryawanshi, K. R., Redpath, S. M., Bhatnagar, Y. V., Ramakrishnan, U., Chaturvedi, V., Smout, S. C., & Mishra, C. (2017). Impact of wild prey availability on livestock predation by snow leopards. *Royal Society open science*, 4(6), 170026.
- Tumursukh, L., Suryawanshi, K. R., Mishra, C., McCarthy, T. M., & Boldgiv, B. (2016). Status of the mountain ungulate prey of the Endangered snow leopard *Panthera uncia* in the Tost Local Protected Area, South Gobi, Mongolia. *Oryx*, 50(2), 214-219.
- Wingard, G. J., Harris, R. B., Amgalanbaatar, S., & Reading, R. P. (2011). Estimating abundance of mountain ungulates incorporating imperfect detection: argali *Ovis ammon* in the Gobi Desert, Mongolia. *Wildlife Biology*, 17(1), 93-101.